F10 Rabbit Polyclonal Antibody





Product Information

CAB15057

Product SKU: Gene ID: CAB15057 2159 20uL, 100uL Size:

Clone No: **Host Species:** Rabbit Reactivity: Human, Mouse, Rat

Additional Information

Observed MW: 56kDa Unconjugated Conjugate:

Calculated MW: 55kDa Isotype: IgG

Immunogen Information

Background: This gene encodes the vitamin K-dependent coagulation factor X of the blood coagulation cascade. This

> factor undergoes multiple processing steps before its preproprotein is converted to a mature two-chain form by the excision of the tripeptide RKR. Two chains of the factor are held together by 1 or more disulfide bonds; the light chain contains 2 EGF-like domains, while the heavy chain contains the catalytic domain which is structurally homologous to those of the other hemostatic serine proteases. The mature factor is activated by the cleavage of the activation peptide by factor IXa (in the intrisic pathway), or by factor VIIa (in the extrinsic pathway). The activated factor then converts prothrombin to thrombin in the presence of factor Va, Ca+2, and phospholipid during blood clotting. Mutations of this gene result in factor X deficiency, a hemorrhagic condition of variable severity. Alternative splicing results in multiple transcript variants encoding different isoforms that may undergo similar proteolytic processing to

generate mature polypeptides.

Recommended Dilution: WB,1:500 - 1:2000 IF/ICC,1:50 - 1:200

Synonyms: FX; FXA; F10

Purifcation Method: Affinity purification

Recombinant fusion protein containing a sequence corresponding to amino acids 41-300 of human F10 Immunogen:

(NP_000495.1).

Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3. Storage: