## **VAV1 Rabbit Polyclonal Antibody**



## **CAB15108**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

110kDa

**Calculated MW:** 

94kDa/98kDa

**Applications:** 

Reactivity:

Human

WB

**Protein Background** 

This gene is a member of the VAV gene family. The VAV proteins are guanine nucleotide exchange factors (GEFs) for Rho family GTPases that activate pathways leading to actin cytoskeletal rearrangements and transcriptional alterations. The encoded protein is important in hematopoiesis, playing a role in T-cell and B-cell development and activation. The encoded protein has been identified as the specific binding partner of Nef proteins from HIV-1. Coexpression and binding of these partners initiates profound morphological changes, cytoskeletal rearrangements and the JNK/SAPK signaling cascade, leading to increased levels of viral transcription and replication. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

Immunogen information

**Gene ID:** 7409

7409

Uniprot P15498

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

Synonyms:

VAV1; VAV

Source:

Immunogen:

Rabbit A synthetic peptide corresponding to a sequence within amino

acids 100-200 of human VAV1 (NP\_005419.2).

Isotype:

lgG Storage:


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of Jurkat cells, using VAV1 antibody (CAB15108) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.