
## **RBM8A Rabbit Polyclonal Antibody**

## CAB15125



Product Information	Protein Background
Size:	This gene encodes a protein with a conserved RNA-binding motif. The protein is found
20uL, 50uL, 100uL, 200uL	predominantly in the nucleus, although it is also present in the cytoplasm. It is preferentially associated with mRNAs produced by splicing, including both nuclear mRNAs and newly
Observed MW:	exported cytoplasmic mRNAs. It is thought that the protein remains associated with spliced mRNAs as a tag to indicate where introns had been present, thus coupling pre- and post-mRNA
20kDa	splicing events. Previously, it was thought that two genes encode this protein, RBM8A and RBM8B; it is now thought that the RBM8B locus is a pseudogene. There are two alternate
Calculated MW:	translation start codons with this gene, which result in two forms of the protein. An allele
19kDa	mutation and a low-frequency noncoding single-nucleotide polymorphism (SNP) in this gene cause thrombocytopenia-absent radius (TAR) syndrome.
Applications:	Immunogen information
WB	
	Gene ID: 9939
Reactivity:	3323
Human, Mouse, Rat	Uniprot Q9Y5S9
Antibody Information	Synonyms:
Recommended dilutions:	RBM8A; BOV-1A; BOV-1B; BOV-1C; C1DELq21.1; DEL1q21.1;
WB 1:500 - 1:2000	MDS014; RBM8; RBM8B; TAR; Y14; ZNRP; ZRNP1
Source:	
Rabbit	Immunogen:
	Recombinant fusion protein containing a sequence corresponding to amino acids 1-80 of human RBM8A (NP_005096.1).
lsotype:	
lgG	Storage:
	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%
	sodium azide, 50% glycerol, pH7.3.
Purification:	
Affinity purification	



Western blot analysis of extracts of various cell lines, using RBM8A antibody (CAB15125) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 10s.