
## ADAMTS1 Rabbit Polyclonal Antibody

## CAB15356



Product Information	Protein Background		
Size:	This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with		
20uL, 50uL, 100uL, 200uL Observed MW: 105kDa Calculated MW: 105kDa Applications:	thrombospondin motif) protein family. Members of the family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the numbe of C-terminal TS motifs, and some have unique C-terminal domains. The protein encoded by this gene contains two disintegrin loops and three C-terminal TS motifs and has anti angiogenic activity. The expression of this gene may be associated with various inflammatory processes as well as development of cancer cachexia. This gene is likely to be necessary fo normal growth, fertility, and organ morphology and function.		
		WB	<b>Gene ID:</b> 9510
		Reactivity:	3510
		Mouse, Rat	<b>Uniprot</b> Q9UHI8
		Antibody Information	Synonyms:
		Beer warde de die die die ee	ADAMTS1; C3-C5; METH1
Recommended dilutions: WB 1:200 - 1:2000			
Source:	Immunogen:		
Rabbit	Recombinant fusion protein containing a sequence corresponding to amino acids 718-967 of human ADAMTS1 (NP_008919.3).		
lsotype:			
lgG	<b>Storage:</b> Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.		

**Purification:** Affinity purification



Western blot analysis of extracts of various cell lines, using ADAMTS1 antibody (CAB15356) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 1s.