
## ACYP2 Rabbit Polyclonal Antibody

## CAB15636



roduct Information	Protein Background
Size:	Acylphosphatase can hydrolyze the phosphoenzyme intermediate of different membrane
20uL, 50uL, 100uL, 200uL	pumps, particularly the Ca2+/Mg2+-ATPase from sarcoplasmic reticulum of skeletal muscle Two isoenzymes have been isolated, called muscle acylphosphatase and erythrocyte
Observed MW:	acylphosphatase on the basis of their tissue localization. This gene encodes the muscle-typ isoform (MT). An increase of the MT isoform is associated with muscle differentiation. Severa
11kDa	transcript variants encoding different isoforms have been found for this gene.
Calculated MW:	Immunogen information
11kDa	Gene ID:
Applications:	98
WB	Uniprot
Reactivity:	P14621
Human, Mouse, Rat	<b>Synonyms:</b> ACYP2; ACYM; ACYP
Antibody Information	
Recommended dilutions:	Immunogen:
WB 1:500 - 1:2000	Recombinant fusion protein containing a sequence corresponding to amino acids 1-99 of human ACYP2 (NP_612457.1).
Source:	
Rabbit	Storage:
	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%
<b>lsotype:</b> lgG	sodium azide, 50% glycerol, pH7.3.

**Purification:** Affinity purification



Western blot analysis of extracts of various cell lines, using ACYP2 antibody (CAB15636) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.