
ENSA Rabbit Polyclonal Antibody

CAB15665



| roduct Information | Protein Background |
|--|--|
| Size: | The protein encoded by this gene belongs to a highly conserved cAMP-regulated |
| 20uL, 50uL, 100uL, 200uL | phosphoprotein (ARPP) family. This protein was identified as an endogenous ligand for the sulfonylurea receptor, ABCC8/SUR1. ABCC8 is the regulatory subunit of the ATP-sensitive |
| Observed MW: | potassium (KATP) channel, which is located on the plasma membrane of pancreatic beta cell and plays a key role in the control of insulin release from pancreatic beta cells. This protein i |
| 13kDa | thought to be an endogenous regulator of KATP channels. In vitro studies have demonstrated that this protein modulates insulin secretion through the interaction with KATP channel, and |
| Calculated MW: | this gene has been proposed as a candidate gene for type 2 diabetes. At least eight alternatively |
| 11kDa/12kDa/13kDa/14kDa/ | spliced transcript variants encoding distinct isoforms have been observed. |
| 15kDa | Immunogen information |
| Applications: | Gene ID: |
| WB | 2029 |
| Reactivity: | Uniprot |
| Mouse | O43768 |
| | Synonyms: |
| Antibody Information | ENSA; ARPP-19e |
| Recommended dilutions: WB 1:500 - 1:2000 | |
| | Immunogen: |
| Source: Rabbit | A synthetic peptide corresponding to a sequence within amino acids 1-100 of human ENSA (NP_996929.1). |
| lsotype: | Storage: |
| IgG | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3. |

Purification: Affinity purification



Western blot analysis of extracts of various cell lines, using ENSA antibody (CAB15665) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 10s.