
AMPKBeta1 Rabbit Polyclonal Antibody

CAB15708



Size:	Protein Background
	The protein encoded by this gene is a regulatory subunit of the AMP-activated protein kinase
20uL, 50uL, 100uL, 200uL	(AMPK). AMPK is a heterotrimer consisting of an alpha catalytic subunit, and non-catalytic beta and gamma subunits. AMPK is an important energy-sensing enzyme that monitors cellula
Observed MW:	energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and inactivates acetyl-CoA carboxylase (ACC) and beta-hydroxy beta
38kDa	methylglutaryl-CoA reductase (HMGCR), key enzymes involved in regulating de novo biosynthesis of fatty acid and cholesterol. This subunit may be a positive regulator of AMPR
Calculated MW:	activity. The myristoylation and phosphorylation of this subunit have been shown to affect the
30kDa	enzyme activity and cellular localization of AMPK. This subunit may also serve as an adapto molecule mediating the association of the AMPK complex.
Applications:	Immunogen information
WB IF	
Reactivity:	Gene ID: 5564
Human	Uniprot Q9Y478
Antibody Information	Synonyms:
Recommended dilutions: WB 1:500 - 1:2000 IF 1:50 - 1:200	PRKAB1; AMPK; HAMPKb
Source:	
Rabbit	Immunogen: A synthetic peptide corresponding to a sequence within amino acids 1-100 of human AMPKBeta1 (NP_006244.2).
lsotype:	
lgG	Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Purification: Affinity purification	



Western blot analysis of extracts of MCF7 cells, using AMPKbeta1 Rabbit pAb (CAB15708) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.

Immunofluorescence analysis of U2OS cells using AMPKbeta1 antibody (CAB15708) at dilution of 1:100. Blue: DAPI for nuclear staining.