## **OSGIN1 Rabbit Polyclonal Antibody**



## **CAB15826**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

75kDa

**Calculated MW:** 

51kDa/60kDa

**Applications:** 

WB

Reactivity:

Human

**Antibody Information** 

Recommended dilutions:

WB 1:500 - 1:2000

Source:

Rabbit

Isotype:

IgG

**Protein Background** 

This gene encodes an oxidative stress response protein that regulates cell death. Expression of the gene is regulated by p53 and is induced by DNA damage. The protein regulates apoptosis by inducing cytochrome c release from mitochondria. It also appears to be a key regulator of both inflammatory and anti-inflammatory molecules. The loss of this protein correlates with uncontrolled cell growth and tumor formation. Naturally occurring read-through transcription exists between this gene and the neighboring upstream malonyl-CoA decarboxylase (MLYCD) gene, but the read-through transcripts are unlikely to produce a protein product.

Immunogen information

**Gene ID:** 29948

23340

**Uniprot** Q9UJX0

**Synonyms:** OSGIN1; BDGI; OKL38

Immunogen:

Recombinant fusion protein containing a sequence corresponding

to amino acids 1-200 of human OSGIN1 (NP\_892026.1).

Storage:


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of 293T cells, using OSGIN1 antibody (CAB15826) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 5s.