
SLC27A4 Rabbit Polyclonal Antibody

CAB16101



roduct Information	Protein Background
Size:	This gene encodes a member of a family of fatty acid transport proteins, which are involved in
20uL, 50uL, 100uL, 200uL	translocation of long-chain fatty acids cross the plasma membrane. This protein is expressed at high levels on the apical side of mature enterocytes in the small intestine, and appears to be
Observed MW:	the principal fatty acid transporter in enterocytes. Clinical studies suggest this gene as a candidate gene for the insulin resistance syndrome. Mutations in this gene have been
72kDa	associated with ichthyosis prematurity syndrome.
Calculated MW:	Immunogen information
26kDa/72kDa	Gene ID:
Applications:	10999
NB	Uniprot
Reactivity:	Q6P1M0
Mouse, Rat	Synonyms:
	SLC27A4; ACSVL4; FATP4; IPS
Antibody Information	
Recommended dilutions:	Immunogen:
WB 1:500 - 1:2000	Recombinant fusion protein containing a sequence corresponding to amino acids 574-643 of human SLC27A4 (NP_005085.2).
Source: Rabbit	
Rubbit	Storage:
	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%
lsotype: IgG	sodium azide, 50% glycerol, pH7.3.

Purification: Affinity purification



Western blot analysis of extracts of various cell lines, using SLC27A4 antibody (CAB16101) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 10s.