PDE11A Rabbit Polyclonal Antibody



CAB16122

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

Refer to figures

Calculated MW:

55kDa/65kDa/78kDa/104kDa

Applications:

WB

Reactivity:

Human, Mouse

Protein Background

The 3', 5'-cyclic nucleotides cAMP and cGMP function as second messengers in a wide variety of signal transduction pathways. 3', 5'-cyclic nucleotide phosphodiesterases (PDEs) catalyze the hydrolysis of cAMP and cGMP to the corresponding 5'-monophosphates and provide a mechanism to downregulate cAMP and cGMP signaling. This gene encodes a member of the PDE protein superfamily. Mutations in this gene are a cause of Cushing disease and adrenocortical hyperplasia. Multiple transcript variants encoding different isoforms have been found for this gene.

Immunogen information

Gene ID: 50940

Uniprot Q9HCR9

Synonyms: PDE11A; PPNAD2

Immunogen:

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000

Source:

Rabbit

Isotype:

IgG

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

Recombinant fusion protein containing a sequence corresponding

to amino acids 712-933 of human PDE11A (NP_058649.3).

sodium azide, 50% glycerol, pH7.3.

Purification:

Affinity purification