## **ALOX12B Rabbit Polyclonal Antibody**



## **CAB16210**

**Product Information** 

20uL, 50uL, 100uL, 200uL

**Protein Background** 

Size:

This gene encodes an enzyme involved in the conversion of arachidonic acid to 12R-

 $hydroxyeicos a tetra enoic\ acid.\ Mutations\ in\ this\ gene\ are\ associated\ with\ nonbullous\ congenital$ 

ichthyosiform erythroderma.

Observed MW:

Immunogen information

80kDa

Gene ID:

Calculated MW: 242

80kDa

Uniprot

**Applications:** O75342

WB

Synonyms:

**Reactivity:** ALOX12B; 12R-LOX; ARCI2; 12R-type

Human, Mouse, Rat

Immunogen:

**Antibody Information** 

Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human ALOX12B (NP\_001130.1).

**Recommended dilutions:** 

WB 1:500 - 1:2000

Storage:

**Source:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

Rabbit sodium azide, 50% glycerol, pH7.3.


Isotype:

IgG

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of Rat testis, using ALOX12B antibody (CAB16210) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 10s.

Western blot analysis of extracts of various cell lines, using ALOX12B antibody (CAB16210) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.