## **NUR77 Rabbit Polyclonal Antibody**



## **CAB16264**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

70kDa

**Calculated MW:** 

33kDa/64kDa/65kDa

**Applications:** 

WB

Reactivity:

Human, Rat

**Protein Background** 

This gene encodes a member of the steroid-thyroid hormone-retinoid receptor superfamily. Expression is induced by phytohemagglutinin in human lymphocytes and by serum stimulation of arrested fibroblasts. The encoded protein acts as a nuclear transcription factor. Translocation of the protein from the nucleus to mitochondria induces apoptosis. Multiple transcript variants encoding different isoforms have been found for this gene.

Immunogen information

**Gene ID:** 3164

**Uniprot** P22736

Synonyms:

NR4A1; GFRP1; HMR; N10; NAK-1; NGFIB; NP10; NUR77; TR3

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

Immunogen:

A synthetic peptide corresponding to a sequence within amino acids 200-300 of human NUR77 (NP\_775180.1).

Source:

Rabbit

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.


Isotype:

IgG

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of rat lung, using NUR77 antibody (CAB16264) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.