

Delta-Catenin/p120 Catenin Rabbit Polyclonal Antibody

CAB1641



Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

108kDa

Calculated MW:

65-72kDa/90-108kDa

Applications:

WB IF

Reactivity:

Human, Mouse

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000 IF 1:50 - 1:200

Source:

Rabbit

Isotype:

IgG

Purification:

Affinity purification

Protein Background

This gene encodes a member of the Armadillo protein family, which function in adhesion between cells and signal transduction. Multiple translation initiation codons and alternative splicing result in many different isoforms being translated. Not all of the full-length natures of the described transcript variants have been determined. Read-through transcription also exists between this gene and the neighboring upstream thioredoxin-related transmembrane protein 2 (TMX2) gene.

Immunogen information

Gene ID:

1500

Uniprot

O60716

Synonyms:

CAS; CTNND; P120CAS; P120CTN; p120; p120(CAS); p120(CTN); CTNND1


Immunogen:

Recombinant fusion protein containing a sequence corresponding to amino acids 573-832 of human Delta-Catenin/p120 Catenin (NP_001078938.1).


Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Product Images



Western blot analysis of extracts of various cell lines, using delta-Catenin/p120 Catenin antibody (CAB1641) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.



Immunofluorescence analysis of U2OS cells using delta-Catenin/p120 Catenin antibody (CAB1641). Blue: DAPI for nuclear staining.