## **CFP Rabbit Polyclonal Antibody**



## **CAB16414**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

51kDa

Calculated MW:

51kDa

**Applications:** 

WB

Reactivity:

Mouse, Rat

-

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000

VVD 1.300 - 1.200

Source:

Rabbit

Isotype:

IgG

Protein Background

This gene encodes a plasma glycoprotein that positively regulates the alternative complement pathway of the innate immune system. This protein binds to many microbial surfaces and apoptotic cells and stabilizes the C3- and C5-convertase enzyme complexes in a feedback loop that ultimately leads to formation of the membrane attack complex and lysis of the target cell. Mutations in this gene result in two forms of properdin deficiency, which results in high susceptibility to meningococcal infections. Multiple alternatively spliced variants, encoding the same protein, have been identified.

Immunogen information

Gene ID:

5199

Uniprot

P27918

Synonyms:

CFP; BFD; PFC; PFD; PROPERDIN; properdin

Immunogen:

Recombinant fusion protein containing a sequence corresponding

to amino acids 190-469 of human CFP (NP\_001138724.1).

Storage:


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of various cell lines, using CFP antibody (CAB16414) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 30s.