
PON2 Rabbit Polyclonal Antibody

CAB1646



roduct Information	Protein Background
Size:	This gene encodes a member of the paraoxonase gene family, which includes three known
20uL, 50uL, 100uL, 200uL	members located adjacent to each other on the long arm of chromosome 7. The encodec protein is ubiquitously expressed in human tissues, membrane-bound, and may act as a cellular
Observed MW:	antioxidant, protecting cells from oxidative stress. Hydrolytic activity against acylhomoserine lactones, important bacterial quorum-sensing mediators, suggests the encoded protein may
37kDa	also play a role in defense responses to pathogenic bacteria. Mutations in this gene may be associated with vascular disease and a number of quantitative phenotypes related to diabetes
Calculated MW:	Alternatively spliced transcript variants encoding different isoforms have been described.
37kDa/39kDa	Immunogen information
Applications:	Gene ID:
WB	5445
Reactivity:	Uniprot
Human	Q15165
	Synonyms:
Antibody Information	PON2
Recommended dilutions: WB 1:500 - 1:2000	
	Immunogen:
Source:	Recombinant fusion protein containing a sequence corresponding
Rabbit	to amino acids 1-210 of human PON2 (NP_000296.2).
lsotype:	Storage:
IgG	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purification: Affinity purification



Western blot analysis of extracts of various cell lines, using PON2 antibody (CAB1646) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.