

# NOMO2 Rabbit Polyclonal Antibody



CAB16614

## Product Information

### Size:

20uL, 50uL, 100uL, 200uL

### Observed MW:

139kDa

### Calculated MW:

122kDa/134kDa/139kDa

### Applications:

WB

### Reactivity:

Human

## Antibody Information

### Recommended dilutions:

WB 1:500 - 1:2000

### Source:

Rabbit

### Isotype:

IgG

### Purification:

Affinity purification

## Protein Background

This gene encodes a protein originally thought to be related to the collagenase gene family. This gene is one of three highly similar genes in a region of duplication located on the p arm of chromosome 16. These three genes encode closely related proteins that may have the same function. The protein encoded by one of these genes has been identified as part of a protein complex that participates in the Nodal signaling pathway during vertebrate development. Mutations in ABCC6, which is located nearby, rather than mutations in this gene are associated with pseudoxanthoma elasticum (PXE). Two transcripts encoding different isoforms have been described.

## Immunogen information

### Gene ID:

283820

### Uniprot

Q5JPE7

### Synonyms:

NOMO2; Nomo; PM5

### Immunogen:


Recombinant fusion protein containing a sequence corresponding to amino acids 873-1222 of human NOMO2 (NP\_001004060.1).

### Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## Product Images

---



Western blot analysis of extracts of BxPC3 cells, using NOMO2 antibody (CAB16614) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 1s.