## **TGFBR1 Rabbit Polyclonal Antibody**



## **CAB16983**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

56kDa

Calculated MW:

56kDa

**Applications:** 

WB IHC IF

Reactivity:

Human, Mouse, Rat

**Antibody Information** 

**Recommended dilutions:** WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Source:

Rabbit

Isotype: IgG

**Purification:** 

Affinity purification

**Protein Background** 

The protein encoded by this gene forms a heteromeric complex with type II TGF-beta receptors when bound to TGF-beta, transducing the TGF-beta signal from the cell surface to the cytoplasm. The encoded protein is a serine/threonine protein kinase. Mutations in this gene have been associated with Loeys-Dietz aortic aneurysm syndrome (LDAS). Multiple transcript variants encoding different isoforms have been found for this gene.

Immunogen information

Gene ID:

7046

Uniprot P36897

Synonyms:

TGFBR1; AAT5; ACVRLK4; ALK-5; ALK5; ESS1; LDS1; LDS1A; LDS2A;

MSSE; SKR4; TGFR-1; tbetaR-I; TBRI; TBR-i

Immunogen:


Recombinant protein of human TGFBR1.


Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

## **Product Images**





Western blot analysis of extracts of various cell lines, using TGFBR1 antibody (CAB16983) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 30s.

Immunofluorescence analysis of NIH/3T3 cells using TGFBR1 antibody (CAB16983) at dilution of 1:100. Blue: DAPI for nuclear staining.