
[KO Validated] ERCC1 Rabbit Polyclonal Antibody

CAB18066



Product Information Size:	Protein Background The product of this gene functions in the nucleotide excision repair pathway, and is required
20uL, 50uL, 100uL, 200uL	for the repair of DNA lesions such as those induced by UV light or formed by electrophilic compounds including cisplatin. The encoded protein forms a heterodimer with the XPF
Observed MW:	endonuclease (also known as ERCC4), and the heterodimeric endonuclease catalyzes the 5' incision in the process of excising the DNA lesion. The heterodimeric endonuclease is also
38kDa	involved in recombinational DNA repair and in the repair of inter-strand crosslinks. Mutations in this gene result in cerebrooculofacioskeletal syndrome, and polymorphisms that alter
Calculated MW:	expression of this gene may play a role in carcinogenesis. Multiple transcript variants encoding
25kDa/29kDa/32kDa/35kDa	different isoforms have been found for this gene. The last exon of this gene overlaps with the CD3e molecule, epsilon associated protein gene on the opposite strand.
Applications:	Immunogen information
WB IF	Gene ID:
Reactivity:	2067
Human, Mouse, Rat	Uniprot P07992
Antibody Information	Synonyms:
Recommended dilutions: WB 1:500 - 1:2000 IF 1:50 - 1:200	ERCC1; COFS4; RAD10; UV20
Source: Rabbit	Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-323 of human ERCC1 (NP_973730.1).
lsotype:	
lgG	Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Purification: Affinity purification	



Western blot analysis of extracts from normal (control) and ERCC1 knockout (KO) HeLa cells, using ERCC1 antibody (CAB18066) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.