

# [KO Validated] ATG4D Rabbit Polyclonal Antibody

## CAB18097



### Product Information

**Size:**

20uL, 50uL, 100uL, 200uL

**Observed MW:**

53kDa

**Calculated MW:**

16kDa/52kDa

**Applications:**

WB IF

**Reactivity:**

Human, Mouse, Rat

### Antibody Information

**Recommended dilutions:**

WB 1:500 - 1:2000 IF 1:50 - 1:200

**Source:**

Rabbit

**Isotype:**

IgG

**Purification:**

Affinity purification

### Protein Background

Autophagy is the process by which endogenous proteins and damaged organelles are destroyed intracellularly. Autophagy is postulated to be essential for cell homeostasis and cell remodeling during differentiation, metamorphosis, non-apoptotic cell death, and aging. Reduced levels of autophagy have been described in some malignant tumors, and a role for autophagy in controlling the unregulated cell growth linked to cancer has been proposed. This gene belongs to the autophagy-related protein 4 (Atg4) family of C54 endopeptidases. Members of this family encode proteins that play a role in the biogenesis of autophagosomes, which sequester the cytosol and organelles for degradation by lysosomes. Alternative splicing results in multiple transcript variants.

### Immunogen information

**Gene ID:**

84971

**Uniprot**

Q86TL0

**Synonyms:**

ATG4D; APG4-D; APG4D; AUTL4

**Immunogen:**


Recombinant fusion protein containing a sequence corresponding to amino acids 1-230 of human ATG4D (NP\_116274.3).

**Storage:**

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## Product Images

---



Western blot analysis of extracts from normal (control) and ATG4D knockout (KO) HeLa cells, using ATG4D antibody (CAB18097) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 3min.