## **PD-L1 Mouse Monoclonal Antibody**



## **CAB18103**

**Product Information** 

Size:

50uL, 100uL, 200uL

**Observed MW:** 

50kDa

**Calculated MW:** 

20kDa/33kDa

**Applications:** 

WB IHC

Reactivity:

Human

**Protein Background** 

This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hematopoietic cells, such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of normal tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer and renal cell carcinoma. Alternative splicing results in multiple transcript variants.

**Immunogen information** 

**Gene ID:** 29126

**Uniprot** Q9NZQ7

Synonyms:

**Antibody Information** 

Recommended dilutions:

WB 1:5000 - 1:10000 IHC

1:50 - 1:200

Source:

Mouse

Mouse

Immunogen:

Recombinant protein of human PD-L1 extracellular domain.

B7-H; B7H1; PDL1; PD-L1; PDCD1L1; PDCD1LG1; CD274

Isotype:

IgG

Storage:


Store at 4°C. Avoid freeze / thaw cycles.Tris Buffer, pH7.3-7.7, with

1% BSA and <0.1% Sodium azide.

**Purification:** 

Affinity purification

## **Product Images**



Western blot - PD-L1 Mouse mAb (CAB18103)