

[KO Validated] Bax Rabbit Monoclonal Antibody



CAB18642

Product Information

Size:

20uL

Observed MW:

21kDa

Calculated MW:

4kDa/12kDa/15kDa/18kDa/19kDa/21kDa/24kDa

Applications:

WB IHC IP

Reactivity:

Human, Mouse

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000 IHC 1:50 - 1:200 IP 1:50 - 1:100

Source:

Rabbit

Isotype:

IgG

Purification:

Affinity purification

Protein Background

The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein forms a heterodimer with BCL2, and functions as an apoptotic activator. This protein is reported to interact with, and increase the opening of, the mitochondrial voltage-dependent anion channel (VDAC), which leads to the loss in membrane potential and the release of cytochrome c. The expression of this gene is regulated by the tumor suppressor P53 and has been shown to be involved in P53-mediated apoptosis. Multiple alternatively spliced transcript variants, which encode different isoforms, have been reported for this gene.

Immunogen information

Gene ID:

581

Uniprot

Q07812

Synonyms:

BCL2L4


Immunogen:

Recombinant protein of human Bax.


Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.


Product Images




Western blot analysis of extracts of various cell lines, using Bax antibody (CAB18642) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 3s.



Western blot analysis of extracts from normal (control) and Bax knockout (KO) 293T cells, using Bax antibody (CAB18642) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 3s.



Immunohistochemistry of paraffin-embedded mouse testis using [KO Validated] Bax Rabbit mAb (CAB18642) at dilution of 1:100 (40x lens).



Immunoprecipitation analysis of 200ug extracts of 293T using 3ug Bax antibody (CAB18642). Western blot was performed from the immunoprecipitate using Bax antibody (CAB18642) at a dilution of 1:1000.