
## TRAF2 Rabbit Monoclonal Antibody

## CAB19129



Product Information	Protein Background
Size:	The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF)
20uL, 50uL, 100uL, 200uL	protein family. TRAF proteins associate with, and mediate the signal transduction from members of the TNF receptor superfamily. This protein directly interacts with TNF receptors,
Observed MW:	and forms a heterodimeric complex with TRAF1. This protein is required for TNF-alpha- mediated activation of MAPK8/JNK and NF-kappaB. The protein complex formed by this
53kDa	protein and TRAF1 interacts with the inhibitor-of-apoptosis proteins (IAPs), and functions as a mediator of the anti-apoptotic signals from TNF receptors. The interaction of this protein with
Calculated MW:	TRADD, a TNF receptor associated apoptotic signal transducer, ensures the recruitment of IAPs for the direct inhibition of caspase activation. BIRC2/c-IAP1, an apoptosis inhibitor possessing
53kDa	ubiquitin ligase activity, can unbiquitinate and induce the degradation of this protein, and thus potentiate TNF-induced apoptosis. Multiple alternatively spliced transcript variants have been
Applications:	found for this gene, but the biological validity of only one transcript has been determined.
WB	[provided by RefSeq, Jul 2008]
Reactivity:	Immunogen information
Human, Mouse, Rat	<b>Gene ID:</b> 7186
Antibody Information	Uniprot Q12933
Recommended dilutions: WB 1:500 - 1:2000 Source: Rabbit	<b>Synonyms:</b> MGC:45012; TRAP; TRAP3; TRAF2
<b>lsotype:</b> lgG	Immunogen: A synthesized peptide derived from human TRAF2
<b>Purification:</b> Affinity purification	<b>Storage:</b> Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot - TRAF2 Rabbit mAb (CAB19129)