
Bad Rabbit Monoclonal Antibody

CAB19595



| Product Information | Protein Background |
|----------------------------|---|
| Size: | The protein encoded by this gene is a member of the BCL-2 family. BCL-2 family members are |
| 20uL, 50uL, 100uL, 200uL | known to be regulators of programmed cell death. This protein positively regulates cel apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor |
| Observed MW: | activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be |
| 23KDa | involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. [provided by RefSeq, Jul 2008] |
| Calculated MW: | transcript variants which encode the same isoform. [provided by Kerseq, Jul 2000] |
| 23kDa | Immunogen information |
| Applications: | Gene ID: 572 |
| WB IHC | |
| Reactivity: | Uniprot Q92934 |
| Human, Mouse, Rat | |
| | Synonyms: BAD; BBC2; BCL2L8; Bad |
| Antibody Information | |
| Recommended dilutions: | |
| WB 1:500 - 1:2000 IHC 1:50 | Immunogen: |
| - 1:200 | A synthesized peptide derived from human Bad. |
| Source: Rabbit | |
| | Storage: |
| | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% |
| lsotype: lgG | sodium azide, 0.05% BSA, 50% glycerol, pH7.3. |
| | |

Purification: Affinity purification



Western blot - Bad Rabbit mAb (CAB19595)