

## CAB19605

### Product Information

<b>Product SKU:</b>	CAB19605	<b>Gene ID:</b>	4791	<b>Size:</b>	20uL, 100uL
<b>Clone No:</b>	ARC0084	<b>Host Species:</b>	Rabbit	<b>Reactivity:</b>	Human,Mouse,Rat

### Additional Information

<b>Observed MW:</b>	120kDa	<b>Conjugate:</b>	Unconjugated
<b>Calculated MW:</b>	97kDa	<b>Isotype:</b>	IgG

### Immunogen Information

<b>Background:</b>	This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFκB). The NFκB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protein encoded by this gene can function as both a transcriptional activator or repressor depending on its dimerization partner. The p100 full-length protein is co-translationally processed into a p52 active form. Chromosomal rearrangements and translocations of this locus have been observed in B cell lymphomas, some of which may result in the formation of fusion proteins. There is a pseudogene for this gene on chromosome 18. Alternative splicing results in multiple transcript variants.
<b>Recommended Dilution:</b>	WB,1:500 - 1:1000 IHC-P,1:50 - 1:200
<b>Synonyms:</b>	p52; p100; H2TF1; LYT10; CVID10; LYT-10; NF-κB2; p49/p100; NF-κB2
<b>Purification Method:</b>	Affinity purification
<b>Immunogen:</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 745-877 of human NF-κB2 (Q00653).
<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.