
Product Information

Product SKU:	CAB19745	Gene ID:	5577	Size:	20uL, 100uL
Clone No:	ARC2272	Host Species:	Rabbit	Reactivity:	Mouse,Rat

Additional Information

Observed MW:	46kDa	Conjugate:	Unconjugated
Calculated MW:	46kDa	Isotype:	IgG

Immunogen Information

Background: cAMP is a signaling molecule important for a variety of cellular functions. cAMP exerts its effects by activating the cAMP-dependent protein kinase, which transduces the signal through phosphorylation of different target proteins. The inactive kinase holoenzyme is a tetramer composed of two regulatory and two catalytic subunits. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits have been identified in humans. The protein encoded by this gene is one of the regulatory subunits. This subunit can be phosphorylated by the activated catalytic subunit. This subunit has been shown to interact with and suppress the transcriptional activity of the cAMP responsive element binding protein 1 (CREB1) in activated T cells. Knockout studies in mice suggest that this subunit may play an important role in regulating energy balance and adiposity. The studies also suggest that this subunit may mediate the gene induction and cataleptic behavior induced by haloperidol.

Recommended Dilution: WB,1:500 - 1:1000 IHC-P,1:50 - 1:200

Synonyms: PRKAR2; RII-BETA; PRKAR2B

Purification Method: Affinity purification

Immunogen: A synthetic peptide corresponding to a sequence within amino acids 319-418 of human PRKAR2B (P31323).

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.