[KO Validated] PMS2 Rabbit Polyclonal Antibody



CAB19927

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

110KDa

Calculated MW:

20kDa/51kDa/62kDa/95kDa

Applications:

WB IF IP

Reactivity:

Human

P54278

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000 IF 1:50 -1:200 IP 1:20 - 1:50

Source:

Rabbit

Isotype:

IgG

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

Affinity purification

Purification:

Protein Background

The protein encoded by this gene is a key component of the mismatch repair system that functions to correct DNA mismatches and small insertions and deletions that can occur during DNA replication and homologous recombination. This protein forms heterodimers with the gene product of the mutL homolog 1 (MLH1) gene to form the MutL-alpha heterodimer. The MutL-alpha heterodimer possesses an endonucleolytic activity that is activated following recognition of mismatches and insertion/deletion loops by the MutS-alpha and MutS-beta heterodimers, and is necessary for removal of the mismatched DNA. There is a DQHA(X)2E(X)4E motif found at the C-terminus of the protein encoded by this gene that forms part of the active site of the nuclease. Mutations in this gene have been associated with hereditary nonpolyposis colorectal cancer (HNPCC; also known as Lynch syndrome) and Turcot syndrome.

Immunogen information

Gene ID:

5395

Uniprot


Synonyms:

Immunogen:

PMS2; HNPCC4; MLH4; PMS2CL; PMSL2

Recombinant protein of human PMS2.

Product Images



Western blot analysis of extracts from normal (control) and PMS2 knockout (KO) HeLa cells, using PMS2 antibody (CAB19927) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 1s.