

**[KO Validated] CDK2AP1 Rabbit Polyclonal
Antibody
CAB19985**



Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

12KDa

Calculated MW:

9kDa/12kDa

Applications:

WB

Reactivity:

Human

Protein Background

The protein encoded by this gene is a cyclin-dependent kinase 2 (CDK2) -associated protein which is thought to negatively regulate CDK2 activity by sequestering monomeric CDK2, and targeting CDK2 for proteolysis. This protein was found to also interact with DNA polymerase alpha/primase and mediate the phosphorylation of the large p180 subunit, which suggests a regulatory role in DNA replication during the S-phase of the cell cycle. This protein also forms a core subunit of the nucleosome remodeling and histone deacetylation (NURD) complex that epigenetically regulates embryonic stem cell differentiation. This gene thus plays a role in both cell-cycle and epigenetic regulation. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Immunogen information

Gene ID:

8099

Uniprot

O14519

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000

Source:

Rabbit

Synonyms:

CDK2AP1; DOC1; DORC1; ST19; doc-1; p12DOC-1

Immunogen:

Recombinant protein of human CDK2AP1.

Isotype:

IgG


Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purification:

Affinity purification

Product Images



Western blot analysis of extracts from normal (control) and CDK2AP1 knockout (KO) 293T cells, using CDK2AP1 antibody (CAB19985) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 30s.