

---

## Product Information

<b>Product SKU:</b>	CAB2011	<b>Gene ID:</b>	51129	<b>Size:</b>	20uL, 100uL
<b>Clone No:</b>	-	<b>Host Species:</b>	Rabbit	<b>Reactivity:</b>	Human,Mouse,Rat

---

## Additional Information

<b>Observed MW:</b>	45-50kDa	<b>Conjugate:</b>	Unconjugated
<b>Calculated MW:</b>	45kDa	<b>Isotype:</b>	IgG

---

## Immunogen Information

<b>Background:</b>	This gene encodes a glycosylated, secreted protein containing a C-terminal fibrinogen domain. The encoded protein is induced by peroxisome proliferation activators and functions as a serum hormone that regulates glucose homeostasis, lipid metabolism, and insulin sensitivity. This protein can also act as an apoptosis survival factor for vascular endothelial cells and can prevent metastasis by inhibiting vascular growth and tumor cell invasion. The C-terminal domain may be proteolytically-cleaved from the full-length secreted protein. Decreased expression of this gene has been associated with type 2 diabetes. Alternative splicing results in multiple transcript variants. This gene was previously referred to as ANGPTL2 but has been renamed ANGPTL4.
<b>Recommended Dilution:</b>	WB,1:500 - 1:1000 IHC-P,1:50 - 1:200 IF/ICC,1:50 - 1:200
<b>Synonyms:</b>	NL2; ARP4; FIAF; HARP; PGAR; HFARP; TGQTL; UNQ171; pp1158; ANGPTL4
<b>Purification Method:</b>	Affinity purification
<b>Immunogen:</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 26-170 of human ANGPTL4 (NP_647475.1).
<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.