## **TGFB1 Rabbit Polyclonal Antibody**





**Product Information** 

**Product SKU**: CAB2124 **Gene ID**: 7040 **Size**: 20uL, 100uL

Clone No: - Host Species: Rabbit Reactivity: Human, Mouse, Rat

**Additional Information** 

**Observed MW**: 12kDa/25kDa/45-60kDa **Conjugate:** Unconjugated

Calculated MW: 44kDa Isotype: IgG

## **Immunogen Information**

**Background**: This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of

proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate a latency-associated peptide (LAP) and a mature peptide, and is found in either a latent form composed of a mature peptide homodimer, a LAP homodimer, and a latent TGF-beta binding protein, or in an active form consisting solely of the mature peptide homodimer. The mature peptide may also form heterodimers with other TGFB family members. This encoded protein regulates cell proliferation, differentiation and growth, and can modulate expression and activation of other growth factors including interferon gamma and tumor necrosis factor alpha. This gene is frequently upregulated in tumor cells, and mutations in this gene result in Camurati-Engelmann disease.

**Recommended Dilution**: WB,1:1000 - 1:5000 IF/ICC,1:50 - 1:200

Synonyms: CED; LAP; DPD1; TGFB; IBDIMDE; TGFbeta; TGF-beta1; TGF beta 1

**Purifcation Method**: Affinity purification

**Immunogen**: A synthetic peptide corresponding to a sequence within amino acids 291-390 of human TGF beta 1

(NP\_000651.3).

**Storage**: Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.