# **Proteasome 20S LMP7 Polyclonal Antibody**



## **CAB22542**

#### **Product Information**

**Product SKU**: CAB22542 **Gene ID**: 5696 **Size**: 20uL, 100uL

Clone No: - Host Species: Rabbit Reactivity: Human, Mouse, Rat

## **Additional Information**

**Observed MW**: 24kDa **Conjugate:** Unconjugated

Calculated MW: 30kDa Isotype: IgG

# **Immunogen Information**

**Background**: The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core

structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. This gene is located in the class II region of the MHC (major histocompatibility complex). Expression of this gene is induced by gamma interferon and this gene product replaces catalytic subunit 3 (proteasome beta 5 subunit) in the immunoproteasome. Proteolytic processing is required to generate a mature subunit. Two alternative transcripts encoding two isoforms have been identified; both isoforms are processed to

yield the same mature subunit.

**Recommended Dilution**: WB,1:500 - 1:1000

Synonyms: JMP; ALDD; LMP7; NKJO; D6S216; PRAAS1; PSMB5i; RING10; D6S216E; Proteasome 20S LMP7

**Purifcation Method**: Affinity purification

**Immunogen**: Recombinant protein of human Proteasome 20S LMP7.

**Storage**: Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.