
## PHD2 Rabbit Polyclonal Antibody

## CAB2314



roduct Information	Protein Background
Size:	The protein encoded by this gene catalyzes the post-translational formation of
20uL, 50uL, 100uL, 200uL	hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. HIF is a transcriptional comple that plays a central role in mammalian oxygen homeostasis. This protein functions as a cellul
Observed MW:	oxygen sensor, and under normal oxygen concentration, modification by prolyl hydroxylatic is a key regulatory event that targets HIF subunits for proteasomal destruction via the vo Hippel-Lindau ubiquitylation complex. Mutations in this gene are associated with erythrocytosis familial type 3 (ECYT3).
Calculated MW:	
36kDa/43kDa/46kDa	Immunogen information
Applications:	<b>Gene ID:</b> 54583
IF	
	Uniprot
Reactivity:	Q9GZT9
Human	
Antibody Information	<b>Synonyms:</b> EGLN1; C1orf12; ECYT3; HALAH; HIF-PH2; HIFPH2; HPH-2; HPH2; PHD2; SM20; ZMYND6
Recommended dilutions:	
IF 1:50 - 1:200	Immunogen:
<b>Source:</b> Rabbit	Recombinant fusion protein containing a sequence corresponding to amino acids 1-426 of human PHD2 (NP_071334.1).
	Storage:
<b>lsotype:</b> lgG	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Purification:** Affinity purification



Immunofluorescence analysis of MCF-7 cells using PHD2 antibody (CAB2314). Blue: DAPI for nuclear staining.