
## 2'-O-methyladenosine / Am Rabbit Polyclonal Antibody

CAB2388



roduct Information	Protein Background		
Size:	Discovered in the 1970s, m6A is the most prevalent internal modification in polyadenylated		
20uL, 50uL, 100uL, 200uL Observed MW: Calculated MW:	mRNAs and long non-coding RNAs (IncRNAs) in higher eukaryotes. m6A is widely conserved among eukaryotic species that range from yeast, plants, flies to mammals, as well as among viral RNAs with a nuclear phase. The m6A-based modification is associated with a well-defined RNA motif, RRACH (R: A/G, H: A/C/U). As a representative of the epitranscriptome, m6A mRNA modifications participate in many vital activities in the cell, including stem cell self-renewal and differentiation, mRNA transcription, alternative splicing, nuclear export, translation degradation, and microRNA processing. These processes determine the expression o inactivation of specific genes, which is vital for growth and development.(PMID: 30416848 PMID: 24662220; PMID: 30429466)		
		Applications:	Immunogen information
		DB	Gene ID:
Reactivity:			
Human, Mouse, Rat, Other (Wide Range)	Uniprot		
Antibody Information	<b>Synonyms:</b> Am; 2'-O-methyladenosine		
Recommended dilutions: DB 1:500 - 1:2000			
Source:	Immunogen: Chemical compounds corresponding to 2'-O-methyladenosine /		
Rabbit	Am.		
lsotype:	-		
IgG	<b>Storage:</b> Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%		
	sodium azide, 50% glycerol, pH7.3.		

Affinity purification



`- 2'-O-methyladenosine / Am Rabbit pAb (CAB2388)