
## **SKP1 Rabbit Polyclonal Antibody**

## CAB2566



Product Information	Protein Background		
<b>Size:</b> 20uL, 50uL, 100uL, 200uL	This gene encodes a component of SCF complexes, which are composed of this protein, cullin 1, a ring-box protein, and one member of the F-box family of proteins. This protein bind directly to the F-box motif found in F-box proteins. SCF complexes are involved in the regulated		
		Observed MW:	ubiquitination of specific protein substrates, which targets them for degradation by th proteosome. Specific F-box proteins recognize different target protein(s), and many specifi
19kDa Calculated MW: 18kDa Applications:	SCF substrates have been identified including regulators of cell cycle progression an development. Studies have also characterized the protein as an RNA polymerase II elongatio factor. Alternative splicing of this gene results in two transcript variants. A related pseudogen has been identified on chromosome 7.		
		WB IHC	<b>Gene ID:</b> 6500
		Reactivity:	
		Human, Mouse	Uniprot P63208
Antibody Information	<b>Synonyms:</b> SKP1; EMC19; OCP-II; OCP2; SKP1A; TCEB1L; p19A		
<b>Recommended dilutions:</b> WB 1:500 - 1:2000 IHC 1:50 - 1:100			
<b>Source:</b> Rabbit	<b>Immunogen:</b> Recombinant fusion protein containing a sequence corresponding to amino acids 1-163 of human SKP1 (NP_733779.1).		
<b>lsotype:</b> lgG	Storage:		
	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.		

**Purification:** Affinity purification



Western blot analysis of extracts of various cell lines, using SKP1 antibody (CAB2566) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 30s.

Immunohistochemistry of paraffin-embedded mouse lung using SKP1 antibody (CAB2566) at dilution of 1:100 (40x lens).