## **LCP2 Rabbit Polyclonal Antibody**



## **CAB2567**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

70kDa

Calculated MW:

60kDa

**Applications:** 

WB

Reactivity:

Human, Mouse

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

Source: Rabbit

Isotype:

IgG

**Purification:** Affinity purification **Protein Background** 

SLP-76 was originally identified as a substrate of the ZAP-70 protein tyrosine kinase following T cell receptor (TCR) ligation in the leukemic T cell line Jurkat. The SLP-76 locus has been localized to human chromosome 5q33 and the gene structure has been partially characterized in mice. The human and murine cDNAs both encode 533 amino acid proteins that are 72% identical and comprised of three modular domains. The NH2-terminus contains an acidic region that includes a PEST domain and several tyrosine residues which are phosphorylated following TCR ligation. SLP-76 also contains a central proline-rich domain and a COOHterminal SH2 domain. A number of additional proteins have been identified that associate with SLP-76 both constitutively and inducibly following receptor ligation, supporting the notion that SLP-76 functions as an adaptor or scaffold protein. Studies using SLP-76 deficient T cell lines or mice have provided strong evidence that SLP-76 plays a positive role in promoting T cell development and activation as well as mast cell and platelet function.

Immunogen information

Gene ID:

3937

Uniprot Q13094

Synonyms:

LCP2; SLP-76; SLP76

Immunogen:

Recombinant fusion protein containing a sequence corresponding


to amino acids 284-533 of human LCP2 (NP\_005556.1).

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

## **Product Images**



Western blot analysis of extracts of various cell lines, using LCP2 antibody (CAB2567) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.