## **TBK1 Rabbit Polyclonal Antibody**



## **CAB2573**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

90kDa

83kDa

**Calculated MW:** 

Applications:

WB

Reactivity:

Mouse

**Protein Background** 

The NF-kappa-B (NFKB) complex of proteins is inhibited by I-kappa-B (IKB) proteins, which inactivate NFKB by trapping it in the cytoplasm. Phosphorylation of serine residues on the IKB proteins by IKB kinases marks them for destruction via the ubiquitination pathway, thereby allowing activation and nuclear translocation of the NFKB complex. The protein encoded by this gene is similar to IKB kinases and can mediate NFKB activation in response to certain growth factors.

Immunogen information

**Gene ID:** 29110

\_\_\_\_\_

**Uniprot** Q9UHD2

**Synonyms:** 

TBK1; FTDALS4; NAK; T2K

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

Immunogen:

Recombinant fusion protein containing a sequence corresponding to amino acids 440-729 of human TBK1 (NP\_037386.1).

Source:

Rabbit

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%


e: sodium azide, 50% glycerol, pH7.3.

**Isotype:** IgG

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of Mouse brain, using TBK1 antibody (CAB2573) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.