

# MonoMethyl-Histone H3-R26 Rabbit Polyclonal Antibody

## CAB3163



### Product Information

**Size:**

20uL, 50uL, 100uL, 200uL

**Observed MW:**

17kDa

**Calculated MW:**

15kDa

**Applications:**

WB IHC IF IP

**Reactivity:**

Human, Mouse, Rat, Other  
(Wide Range)

### Antibody Information

**Recommended dilutions:**

WB 1:500 - 1:2000 IHC 1:50  
- 1:200 IF 1:50 - 1:200 IP  
1:50 - 1:100

**Source:**

Rabbit

**Isotype:**

IgG

**Purification:**

Affinity purification

### Protein Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

### Immunogen information

**Gene ID:**

8290

**Uniprot**

Q16695

**Synonyms:**

H3.4; H3/g; H3FT; H3t; HIST3H3; Histone H3; HIST1H3A


**Immunogen:**

A synthetic methylated peptide corresponding to residues surrounding R26 of human histone H3


**Storage:**

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


## Product Images




Western blot analysis of extracts of various cell lines, using MonoMethyl-Histone H3-R26 antibody (CAB3163). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.



Dot-blot analysis of all sorts of methylation peptides using MonoMethyl-Histone H3-R26 antibody (CAB3163).



Immunohistochemistry of paraffin-embedded mouse testis using H3R26me1 antibody (CAB3163) at dilution of 1:100 (40x lens).



Immunofluorescence analysis of 293T cells using MonoMethyl-Histone H3-R26 antibody (CAB3163). Blue: DAPI for nuclear staining.