[KO Validated] GSK3Beta Rabbit Polyclonal **Antibody**



CAB3174

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

46kDa

Calculated MW:

46kDa/48kDa

Applications:

WB

Reactivity:

Human, Rat

Antibody Information Recommended dilutions:

WB 1:500 - 1:1000

Source: Rabbit

Isotype: IgG

Protein Background

The protein encoded by this gene is a serine-threonine kinase, belonging to the glycogen synthase kinase subfamily. It is involved in energy metabolism, neuronal cell development, and body pattern formation. Polymorphisms in this gene have been implicated in modifying risk of Parkinson disease, and studies in mice show that overexpression of this gene may be relevant to the pathogenesis of Alzheimer disease. Alternatively spliced transcript variants encoding

different isoforms have been found for this gene.

Immunogen information

Gene ID: 2932

Uniprot P49841

Synonyms: GSK3B; qsk-3Beta

Immunogen:

A synthetic peptide corresponding to a sequence within amino

acids 300 to the C-terminus of human GSK3Beta

(NP_001139628.1).


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

Purification:

Affinity purification

Product Images



Western blot analysis of extracts of various cell lines, using GSK3beta antibody (CAB3174) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.

Western blot analysis of extracts from normal (control) and GSK3beta knockout (KO) HeLa cells, using GSK3b antibody (CAB3174) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 30s.