[KO Validated] SNF5 Rabbit Monoclonal Antibody



CAB3247

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

44KDa

44kDa

Calculated MW:

Applications:

WB IF

Reactivity:

Human, Mouse, Rat

Antibody Information

Recommended dilutions: WB 1:500 - 1:2000 IF 1:50 -

1:200

Source:

Rabbit

Isotype: IgG

Purification:

Affinity purification

Protein Background

The protein encoded by this gene is part of a complex that relieves repressive chromatin structures, allowing the transcriptional machinery to access its targets more effectively. The encoded nuclear protein may also bind to and enhance the DNA joining activity of HIV-1 integrase. This gene has been found to be a tumor suppressor, and mutations in it have been associated with malignant rhabdoid tumors. Alternatively spliced transcript variants have been

found for this gene. [provided by RefSeq, Dec 2015]

Immunogen information

Gene ID:

6598

Uniprot

Q12824

Synonyms:

BAF47; CSS3; INI1; MRD15; PPP1R144; RDT; RTPS1; SNF5; SNF5L1;

SWNTS1; Sfh1p; Snr1; hSNFS

Immunogen:


A synthesized peptide derived from human SNF5

Storage:


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 0.05% BSA, 50% glycerol, pH7.3.


Product Images




Western blot analysis of extracts of various cell lines, using SNF5 Rabbit mAb (CAB3247) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 3s.



Western blot analysis of extracts of various cell lines, using SNF5 Rabbit mAb (CAB3247) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 10s.



Western blot analysis of extracts of 293T cells, using SNF5 antibody (CAB3247) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 10s.



Immunofluorescence analysis of NIH-3T3 cells using SNF5 Rabbit mAb (CAB3247) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.