## **TSTA3 Rabbit Polyclonal Antibody**



## **CAB4169**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

36kDa

35kDa

WB

Calculated MW:

Applications:

Reactivity:

Human, Mouse

Recommended dilutions:

**Antibody Information** 

WB 1:1000 - 1:2000

Source:

Rabbit

**Isotype:** IgG

**Purification:** 

Affinity purification

**Protein Background** 

Tissue specific transplantation antigen P35B is a NADP(H)-binding protein. It catalyze the two-step epimerase and the reductase reactions in GDP-D-mannose metabolism, converting GDP-4-keto-6-D-deoxymannose to GDP-L-fucose. GDP-L-fucose is the substrate of several fucosyltransferases involved in the expression of many glycoconjugates, including blood group ABH antigens and developmental adhesion antigens. Mutations in this gene may cause

leukocyte adhesion deficiency, type II.

Immunogen information

Gene ID:

7264

Uniprot

Q13630

**Synonyms:** 

TSTA3; FX; P35B; SDR4E1

Immunogen:

Recombinant fusion protein containing a sequence corresponding


to amino acids 1-321 of human TSTA3 (NP\_003304.1).

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

## **Product Images**



Western blot analysis of extracts of various cell lines, using TSTA3 antibody (CAB4169) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 10s.