
## JUNB Rabbit Polyclonal Antibody

## CAB5290




| Product Information  | Protein Background   |
|--|--|
| Size:  | JunB is one of the components of the Activator Protein-1 (AP-1) transcription complex that   |
| 20uL, 50uL, 100uL, 200uL   | have been implicated in the control of the G0/G1 transiton in fibroblasts. AP-1 is a collection or<br>dimers formed by members of the Jun-, Fos-, ATF- and Maf multigene families that bind to |
| Observed MW:   | specific DNA regulatory elements called AP-1/12-O-tetradecanoylphorbol-13-acetate-<br>responsive elements (TREs) and cAMP-responsive elements (CREs). JunB binds to the DNA                    |
| 43KDa  | sequence 5'-TGA[CG]TCA-3', and involves in the regulation of gene activity following the primary growth factor response. It also acts either as a tumor suppressor or as an oncogene           |
| Calculated MW:   | depending on the cell and physiopathological context   |
| 35kDa  | Immunogen information  |
| Applications:  | Gene ID:   |
| WB IHC IF  | 3726   |
| Reactivity:  | Uniprot  |
| Human, Mouse, Rat  | P17275   |
|  | Synonyms:  |
| Antibody Information   | JUNB; AP-1   |
| <b>Recommended dilutions:</b><br>WB 1:500 - 1:2000 IHC 1:50<br>- 1:200 IF 1:50 - 1:100 | Immunogen:   |
| Source:  | Recombinant fusion protein containing a sequence corresponding   |
| Rabbit   | to amino acids 1-240 of human JUNB (NP_002220.1).  |
| lsotype:   | Storage:   |
| lgG  | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.  |

**Purification:** Affinity purification



Western blot analysis of extracts of Mouse liver, using JUNB antibody (CAB5290) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 1s.



Immunofluorescence analysis of HeLa cells using JUNB antibody (CAB5290).