
TAF12 Rabbit Polyclonal Antibody

CAB5421



Product Information	Protein Background
Size:	Control of transcription by RNA polymerase II involves the basal transcription machinery which
20uL, 50uL, 100uL, 200uL	is a collection of proteins. These proteins with RNA polymerase II, assemble into complexes which are modulated by transactivator proteins that bind to cis-regulatory elements located
Observed MW:	adjacent to the transcription start site. Some modulators interact directly with the basal complex, whereas others may act as bridging proteins linking transactivators to the basal
20kDa	transcription factors. Some of these associated factors are weakly attached while others are tightly associated with TBP in the TFIID complex. Among the latter are the TAF proteins.
Calculated MW:	Different TAFs are predicted to mediate the function of distinct transcriptional activators for a
14kDa/17kDa	variety of gene promoters and RNA polymerases. TAF12 interacts directly with TBP as well as with TAF2I. Two transcript variants encoding the same protein have been found for this gene.
Applications:	Immunogen information
WB	5
	Gene ID:
Reactivity:	6883
Human	Uniprot Q16514
Antibody Information	Synonyms:
Recommended dilutions: WB 1:500 - 1:2000	TAF12; TAF2J; TAFII20
Source:	
Rabbit	Immunogen:
	Recombinant fusion protein containing a sequence corresponding to amino acids 1-161 of human TAF12 (NP_001128690.1).
lsotype:	
lgG	Storage
	Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Purification:	
Affinity purification	



Western blot analysis of extracts of various cell lines, using TAF12 antibody (CAB5421) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (CABM00021). Exposure time: 120s.