

**CAB5476**

## Product Information

<b>Product SKU:</b>	CAB5476	<b>Gene ID:</b>	5824	<b>Size:</b>	20uL, 100uL
<b>Clone No:</b>	-	<b>Host Species:</b>	Rabbit	<b>Reactivity:</b>	Human,Mouse,Rat

## Additional Information

<b>Observed MW:</b>	37kDa	<b>Conjugate:</b>	Unconjugated
<b>Calculated MW:</b>	33kDa	<b>Isotype:</b>	IgG

## Immunogen Information

<b>Background:</b>	This gene is necessary for early peroxisomal biogenesis. It acts both as a cytosolic chaperone and as an import receptor for peroxisomal membrane proteins (PMPs). Peroxisins (PEXs) are proteins that are essential for the assembly of functional peroxisomes. The peroxisome biogenesis disorders (PBDs) are a group of genetically heterogeneous autosomal recessive, lethal diseases characterized by multiple defects in peroxisome function. These disorders have at least 14 complementation groups, with more than one phenotype being observed for some complementation groups. Although the clinical features of PBD patients vary, cells from all PBD patients exhibit a defect in the import of one or more classes of peroxisomal matrix proteins into the organelle. Defects in this gene are a cause of Zellweger syndrome (ZWS), as well as peroxisome biogenesis disorder complementation group 14 (PBD-CG14), which is also known as PBD-CGJ. Alternative splicing results in multiple transcript variants.
<b>Recommended Dilution:</b>	WB,1:500 - 1:2000 IF/ICC,1:50 - 1:200
<b>Synonyms:</b>	PXF; HK33; PMP1; PMPI; PXMP1; PBD12A; D1S2223E; PEX19
<b>Purification Method:</b>	Affinity purification
<b>Immunogen:</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 1-299 of human PEX19 (NP_002848.1).
<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.