
## F2R Rabbit Polyclonal Antibody

## CAB5641



Product Information Size: 20uL, 50uL, 100uL, 200uL Observed MW:	<b>Protein Background</b> Coagulation factor II receptor is a 7-transmembrane receptor involved in the regulation of thrombotic response. Proteolytic cleavage leads to the activation of the receptor. F2R is a G- protein coupled receptor family member. Alternative splicing results in multiple transcript variants.		
		47kDa	Immunogen information
		Calculated MW:	Gene ID:
		47kDa	2149
Applications:	Uniprot		
WB	P25116		
Reactivity:	Synonyms:		
Human, Mouse, Rat	F2R; CF2R; HTR; PAR-1; PAR1; TR		
Antibody Information	Immunogen:		
Recommended dilutions: WB 1:500 - 1:2000	Recombinant fusion protein containing a sequence corresponding to amino acids 42-110 of human F2R (NP_001983.2).		
Source:	Storage:		
Rabbit	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.		
lsotype:			
lgG			

**Purification:** Affinity purification



Western blot analysis of extracts of various cell lines, using F2R antibody (CAB5641) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 5s.