
## EHHADH Rabbit Polyclonal Antibody

## CAB5717



roduct Information	Protein Background		
Size:	The protein encoded by this gene is a bifunctional enzyme and is one of the four enzymes o		
20uL, 50uL, 100uL, 200uL <b>Observed MW:</b> 79kDa	the peroxisomal beta-oxidation pathway. The N-terminal region of the encoded protein contains enoyl-CoA hydratase activity while the C-terminal region contains 3-hydroxyacyl-CoA dehydrogenase activity. Defects in this gene are a cause of peroxisomal disorders such as Zellweger syndrome. Two transcript variants encoding different isoforms have been found fo this gene.		
		Calculated MW:	Immunogen information
		69kDa/79kDa	Gene ID:
Applications:	1962		
WB	Uniprot		
Reactivity:	Q08426		
Human, Mouse, Rat	<b>Synonyms:</b> EHHADH; ECHD; FRTS3; L-PBE; LBFP; LBP; PBFE		
Antibody Information			
Recommended dilutions: WB 1:500 - 1:2000	<b>Immunogen:</b> Recombinant fusion protein containing a sequence corresponding to amino acids 444-723 of human EHHADH (NP_001957.2).		
<b>Source:</b> Rabbit			
	Storage:		
<b>lsotype:</b> lgG	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.		

**Purification:** Affinity purification



Western blot analysis of extracts of various cell lines, using EHHADH antibody (CAB5717) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.