
QDPR Rabbit Polyclonal Antibody

CAB5733



Product Information	Protein Background
Size:	This gene encodes the enzyme dihydropteridine reductase, which catalyzes the NADH
20uL, 50uL, 100uL, 200uL	mediated reduction of quinonoid dihydrobiopterin. This enzyme is an essential component o the pterin-dependent aromatic amino acid hydroxylating systems. Mutations in this gene
Observed MW:	resulting in QDPR deficiency include aberrant splicing, amino acid substitutions, insertions, o premature terminations. Dihydropteridine reductase deficiency presents as atypica
26kDa	phenylketonuria due to insufficient production of biopterin, a cofactor for phenylalaning hydroxylase.
Calculated MW:	
22kDa/25kDa	Immunogen information
Applications:	Gene ID:
	5860
WB IF	
Reactivity:	Uniprot P09417
Human, Mouse, Rat	
	Synonyms: QDPR; DHPR; PKU2; SDR33C1
Antibody Information	
Recommended dilutions:	
WB 1:500 - 1:2000 IF 1:50 -	Immunogen:
1:100	Recombinant fusion protein containing a sequence corresponding
Source: Rabbit	to amino acids 1-244 of human QDPR (NP_000311.2).
	Storage:
lsotype:	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%
lgG	sodium azide, 50% glycerol, pH7.3.

Purification: Affinity purification



Western blot analysis of extracts of various cell lines, using QDPR antibody (CAB5733) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.

Western blot analysis of extracts of various cell lines, using QDPR antibody (CAB5733) at 1:3000 dilution._Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution._Lysates/proteins: 25ug per lane._Blocking buffer: 3% nonfat dry milk in TBST._Detection: ECL Enhanced Kit (CABM00021)._Exposure time: 90s.

Immunofluorescence analysis of U2OS cells using QDPR antibody (CAB5733). Blue: DAPI for nuclear staining.

