
IL2RB Rabbit Polyclonal Antibody

CAB6207



The interleukin 2 receptor, which is involved in T cell-mediated immune responses, is preser in 3 forms with respect to ability to bind interleukin 2. The low affinity form is a monomer of the alpha subunit and is not involved in signal transduction. The intermediate affinity form consists of an alpha/beta subunit heterodimer, while the high affinity form consists of a
the alpha subunit and is not involved in signal transduction. The intermediate affinity form
consists of an alpha/beta subunit heterodimer, while the high affinity form consists of a
alpha/beta/gamma subunit heterotrimer. Both the intermediate and high affinity forms of th
receptor are involved in receptor-mediated endocytosis and transduction of mitogenic signal from interleukin 2. The protein encoded by this gene represents the beta subunit and is a typ
I membrane protein.
Immunogen information
Gene ID:
3560
Uniprot
P14784
Synonyms:
IL2RB; CD122; IL15RB; P70-75
Immunogen:
Recombinant fusion protein containing a sequence corresponding to amino acids 27-240 of human IL2RB (NP_000869.1).
Storage:
Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purification: Affinity purification



Western blot analysis of extracts of various cell lines, using IL2RB antibody (CAB6207) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 10s.

Western blot analysis of extracts of RAW264.7 cells, using IL2RB antibody (CAB6207) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 30s.