UCN2 Rabbit Polyclonal Antibody

CAB6485




| Product Information | Protein Background |
|--|--|
| Size: | This gene is a member of the sauvagine/corticotropin-releasing factor/urotensin I family. It i |
| 20uL, 50uL, 100uL, 200uL | structurally related to the corticotropin-releasing factor (CRF) gene and the encoded product is an endogenous ligand for CRF type 2 receptors. In the brain it may be responsible for the effects of stress on appetite. In spite of the gene family name similarity, the product of this gene has no sequence similarity to urotensin II. |
| Observed MW: | |
| 12kDa | |
| Calculated MW: | Immunogen information |
| 12kDa | Gene ID: 90226 |
| Applications: | |
| WB IHC IF | Uniprot Q96RP3 |
| Reactivity: | |
| Human, Mouse, Rat | Synonyms: UCN2; SRP; UCN-II; UCNI; UR; URP |
| Antibody Information | |
| Recommended dilutions: WB 1:500 - 1:2000 IHC 1:50 - 1:100 IF 1:50 - 1:100 | Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 20-112 of human UCN2 (NP_149976.1). |
| Source: | Storage |
| Rabbit | Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3. |

lsotype: lgG

Purification: Affinity purification

Product Images



Western blot analysis of extracts of rat skin, using UCN2 antibody (CAB6485) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.

Immunohistochemistry of paraffin-embedded mouse heart using UCN2 antibody (CAB6485) at dilution of 1:100 (40x lens).

Immunofluorescence analysis of U2OS cells using UCN2 antibody (CAB6485). Blue: DAPI for nuclear staining.