ATP6V1B1 Rabbit Polyclonal Antibody

CAB6876

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

57kDa

Calculated MW:

56kDa

Applications:

WB

Reactivity:

Human, Mouse

Antibody Information

Recommended dilutions:
WB 1:500 - 1:2000

Source:
Rabbit

Isotype:
IgG

Purification:
Affinity purification

Protein Background

This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is one of two V1 domain B subunit isoforms and is found in the kidney. Mutations in this gene cause distal renal tubular acidosis associated with sensorineural deafness.

Immunogen information

Gene ID:
525

Uniprot
P15313

Synonyms:
ATP6V1B1; ATP6B1; RTA1B; VATB; VMA2; VPP3

Immunogen:
Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human ATP6V1B1 (NP_001683.2).

Storage:
Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Product Images

Western blot analysis of extracts of various cell lines, using ATP6V1B1 antibody (CAB6876) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.