GBP1 Rabbit Polyclonal Antibody



CAB6911

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

68kDa

Calculated MW:

Applications:

67kDa

WB IF

Reactivity:

Human

Protein Background

Guanylate binding protein expression is induced by interferon. Guanylate binding proteins are characterized by their ability to specifically bind guanine nucleotides (GMP, GDP, and GTP) and are distinguished from the GTP-binding proteins by the presence of 2 binding motifs rather

than 3.

Immunogen information

Gene ID: 2633

Uniprot P32455

Synonyms:

Immunogen:

GBP1

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000 IF 1:50 - 1:100

Source:

Source

Rabbit

Isotype:

lgG

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

Recombinant fusion protein containing a sequence corresponding


to amino acids 383-592 of human GBP1 (NP_002044.2).


sodium azide, 50% glycerol, pH7.3.

Purification:

Affinity purification

Product Images





Western blot analysis of extracts of 5637(HTB-9) cells, using GBP1 antibody (CAB6911) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 10s.

Immunofluorescence analysis of U2OS cells using GBP1 antibody (CAB6911). Blue: DAPI for nuclear staining.