# TriMethyl-Histone H3-K56 Rabbit Polyclonal Antibody

CAB7262



#### Product Information Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

16kDa

Calculated MW:

15kDa

#### Applications:

WB IHC IF IP ChIP ChIPseq

#### **Reactivity:**

Human, Mouse, Rat, Other (Wide Range)

## **Antibody Information**

#### **Recommended dilutions:**

WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200 IP 1:50 - 1:200 ChIP 1:20 -1:100 ChIPseq 1:20 - 1:100 **Source:** Rabbit

**lsotype:** lgG

**Purification:** Affinity purification

#### **Protein Background**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

### Immunogen information

Gene ID: 8290

**Uniprot** Q16695

#### Synonyms:


H3.4; H3/g; H3FT; H3t; HIST3H3; Histone H3; HIST1H3A

Immunogen:


A synthetic peptide of human TriMethyl-Histone H3-K56

#### Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.



Western blot analysis of extracts of various cell lines, using TriMethyl-Histone H3-K56 antibody (CAB7262) at 1:1000 dilution.\_Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution.\_Lysates/proteins: 25ug per lane.\_Blocking buffer: 3% nonfat dry milk in TBST.\_Detection: ECL Enhanced Kit (CABM00021).\_Exposure time: 90s.



Immunofluorescence analysis of HeLa cells using TriMethyl-Histone H3-K56 pAb (CAB7262) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.